

# The Theoretical Minimum

## Quantum Mechanics - Solutions

L02E02

Last version: [tales.mbivert.com/on-the-theoretical-minimum-solutions/](https://tales.mbivert.com/on-the-theoretical-minimum-solutions/) or [github.com/mbivert/ttm](https://github.com/mbivert/ttm)

M. Bivert

June 23, 2023

**Exercise 1.** Prove that  $|i\rangle$  and  $|o\rangle$  satisfy all of the conditions in Eqs. 2.7, 2.8 and 2.9. Are they unique in that respect?

---

Let us recall, in order, Eqs. 2.7, 2.8, 2.9, 2.10, which defines  $|i\rangle$  and  $|o\rangle$ , and both 2.5 and 2.6 which defines  $|r\rangle$  and  $|l\rangle$ :

$$\langle i|o\rangle = 0$$

$$\begin{aligned}\langle o|u\rangle \langle u|o\rangle &= \frac{1}{2} \\ \langle i|u\rangle \langle u|i\rangle &= \frac{1}{2}\end{aligned}$$

$$\begin{aligned}\langle o|d\rangle \langle d|o\rangle &= \frac{1}{2} \\ \langle i|d\rangle \langle d|i\rangle &= \frac{1}{2}\end{aligned}$$

$$\begin{aligned}\langle o|r\rangle \langle r|o\rangle &= \frac{1}{2} \\ \langle i|r\rangle \langle r|i\rangle &= \frac{1}{2}\end{aligned}$$

$$\begin{aligned}\langle o|l\rangle \langle l|o\rangle &= \frac{1}{2} \\ \langle i|l\rangle \langle l|i\rangle &= \frac{1}{2}\end{aligned}$$

$$|i\rangle = \frac{1}{\sqrt{2}}|u\rangle + \frac{i}{\sqrt{2}}|d\rangle$$

$$|o\rangle = \frac{1}{\sqrt{2}}|u\rangle - \frac{i}{\sqrt{2}}|d\rangle$$

$$|r\rangle = \frac{1}{\sqrt{2}}|u\rangle + \frac{1}{\sqrt{2}}|d\rangle$$

$$|l\rangle = \frac{1}{\sqrt{2}}|u\rangle - \frac{1}{\sqrt{2}}|d\rangle$$

For clarity, let us recall that  $\langle u|A\rangle$  is the component of  $|A\rangle$  along the orthonormal vector  $|u\rangle$ . This is because in an *orthonormal* basis  $(|i\rangle)_{i \in F}$  we have:

$$\begin{aligned}|A\rangle &= \sum_{i \in F} \alpha_i |i\rangle \\ \Rightarrow \langle j|A\rangle &= \langle j| \sum_{i \in F} \alpha_i |i\rangle = \sum_{i \in F} \alpha_i \underbrace{\langle j|i\rangle}_{=\delta_{ij}} = \alpha_j\end{aligned}$$

And to make better sense of those equations, let us recall that  $\alpha_u^* \alpha_u = \langle A|u \rangle \langle u|A \rangle$  is the probability of a state vector  $|A\rangle = \alpha_u|u\rangle + \alpha_d|d\rangle$  to be measured in the state  $|u\rangle$ .

For Eq. 2.7, we have

$$\begin{aligned} \langle i|o\rangle &= (\iota_u^* \ \iota_d^*) \begin{pmatrix} o_u \\ o_d \end{pmatrix} \\ &= \iota_u^* o_u + \iota_d^* o_d \\ &= \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} + \frac{-i}{\sqrt{2}} \frac{-i}{\sqrt{2}} = \frac{1}{2} - \frac{1}{2} = 0 \quad \square \end{aligned}$$

For Eqs. 2.8, we can rely on the projection on an orthonormal vector:

$$\begin{aligned} \langle o|u\rangle \langle u|o\rangle &= \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} = \frac{1}{2} \quad \square & \langle o|d\rangle \langle d|o\rangle &= \frac{i}{\sqrt{2}} \frac{-i}{\sqrt{2}} = \frac{1}{2} \quad \square \\ \langle i|u\rangle \langle u|i\rangle &= \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} = \frac{1}{2} \quad \square & \langle i|d\rangle \langle d|i\rangle &= \frac{-i}{\sqrt{2}} \frac{i}{\sqrt{2}} = \frac{1}{2} \quad \square \end{aligned}$$

For Eqs. 2.9, we need to rely on the column form of the inner-product:

$$\begin{aligned} \langle o|r\rangle \langle r|o\rangle &= (o_u^* \ o_d^*) \begin{pmatrix} \rho_u \\ \rho_d \end{pmatrix} (\rho_u^* \ \rho_d^*) \begin{pmatrix} o_u \\ o_d \end{pmatrix} & \langle o|l\rangle \langle l|o\rangle &= (o_u^* \ o_d^*) \begin{pmatrix} \lambda_u \\ \lambda_d \end{pmatrix} (\lambda_u^* \ \lambda_d^*) \begin{pmatrix} o_u \\ o_d \end{pmatrix} \\ &= \left( \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}} \frac{1}{\sqrt{2}} \right) \left( \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} \frac{-i}{\sqrt{2}} \right) & &= \left( \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} + \frac{i}{\sqrt{2}} \frac{-1}{\sqrt{2}} \right) \left( \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} + \frac{-1}{\sqrt{2}} \frac{-i}{\sqrt{2}} \right) \\ &= \left( \frac{1}{2} + \frac{i}{2} \right) \left( \frac{1}{2} - \frac{i}{2} \right) & &= \left( \frac{1}{2} - \frac{i}{2} \right) \left( \frac{1}{2} + \frac{i}{2} \right) \\ &= \frac{1}{4} (1+i)(1-i) & &= \frac{1}{4} (1-i)(1+i) \\ &= \frac{1}{4} (1+i-i+1) = \frac{1}{2} \quad \square & &= \frac{1}{4} (1-i+i+1) = \frac{1}{2} \quad \square \\ \langle i|r\rangle \langle r|i\rangle &= (\iota_u^* \ \iota_d^*) \begin{pmatrix} \rho_u \\ \rho_d \end{pmatrix} (\rho_u^* \ \rho_d^*) \begin{pmatrix} \iota_u \\ \iota_d \end{pmatrix} & \langle i|l\rangle \langle l|i\rangle &= (\iota_u^* \ \iota_d^*) \begin{pmatrix} \lambda_u \\ \lambda_d \end{pmatrix} (\lambda_u^* \ \lambda_d^*) \begin{pmatrix} \iota_u \\ \iota_d \end{pmatrix} \\ &= \left( \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} + \frac{-i}{\sqrt{2}} \frac{1}{\sqrt{2}} \right) \left( \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} \frac{i}{\sqrt{2}} \right) & &= \left( \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} + \frac{-i}{\sqrt{2}} \frac{-1}{\sqrt{2}} \right) \left( \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} + \frac{-1}{\sqrt{2}} \frac{i}{\sqrt{2}} \right) \\ &= \left( \frac{1}{2} - \frac{i}{2} \right) \left( \frac{1}{2} + \frac{i}{2} \right) & &= \left( \frac{1}{2} + \frac{i}{2} \right) \left( \frac{1}{2} - \frac{i}{2} \right) \\ &= \frac{1}{4} (1-i)(1+i) & &= \frac{1}{4} (1+i)(1-i) \\ &= \frac{1}{4} (1+i+i+1) = \frac{1}{2} \quad \square & &= \frac{1}{4} (1+i-i+1) = \frac{1}{2} \quad \square \end{aligned}$$

---

Regarding the unicity of  $|i\rangle, |o\rangle$ , as for  $|r\rangle, |l\rangle$ , there definitely is a phase ambiguity, meaning, we can multiply either  $|i\rangle$  or  $|o\rangle$  by a *phase factor*, say  $e^{i\theta}$ , without disturbing any of the constraints: orthogonality, probabilities, and the resulting vectors are still unitary.

But as stated by the authors for  $|r\rangle, |l\rangle$ , measurable quantities are independant of any phase factors. Thus, so far, there seems to be unicity, up to such a phase factor.

**Remark 1.** *I think some sort of dimensional argument might be required to rigorously prove that indeed there's no way to extract more than three pairs of mutually orthogonal vectors which have a inner-product to 1/2, in a  $\mathbb{C}$ -vector space setting.*