

# The Theoretical Minimum

## Quantum Mechanics - Solutions

L04E05

Last version: [tales.mbivert.com/on-the-theoretical-minimum-solutions/](https://tales.mbivert.com/on-the-theoretical-minimum-solutions/) or [github.com/mbivert/ttm](https://github.com/mbivert/ttm)

M. Bivert

May 10, 2023

**Exercise 1.** Take any unit 3-vector  $\mathbf{n}$  and form the operator

$$H = \frac{\hbar\omega}{2} \boldsymbol{\sigma} \cdot \mathbf{n}$$

Find the energy eigenvalues and eigenvectors by solving the time-independent Schrödinger equation. Recall that Eq. 3.23 gives  $\boldsymbol{\sigma} \cdot \mathbf{n}$  in component form.

Let's recall Eq. 3.23, which is general form of the spin 3-vector operator:

$$\sigma_n = \boldsymbol{\sigma} \cdot \mathbf{n} = \begin{pmatrix} n_z & (n_x - in_y) \\ (n_x + in_y) & -n_z \end{pmatrix}$$

And the time-independent Schrödinger equation<sup>1</sup>:

$$H|E_j\rangle = E_j|E_j\rangle$$

In an earlier exercise (L03E04), we actually diagonalized  $\sigma_n$ : this gave us two eigenvalues  $+1$  and  $-1$ , and two eigenvectors:

$$|+1\rangle = \begin{pmatrix} \cos(\theta/2) \\ \exp(i\phi) \sin(\theta/2) \end{pmatrix}; \quad |-1\rangle = \begin{pmatrix} -\sin(\theta/2) \\ \exp(i\phi) \cos(\theta/2) \end{pmatrix}$$

Where  $\mathbf{n}$  was a regular unitary 3-vector expressed in spherical coordinates:

$$\mathbf{n} = \begin{pmatrix} \sin \theta \cos \phi \\ \sin \theta \sin \phi \\ \cos \theta \end{pmatrix}$$

Let's see how we can leverage this previous work to our advantage: such an  $\mathbf{n}$  vector still fit our purpose here. Furthermore, we know that the eigenvalues of  $\sigma_n$  are the only solutions to:

$$\sigma_n|F_j\rangle = F_j|F_j\rangle$$

But if we multiply both sides of this equation by  $\frac{\hbar\omega}{2}$ , we get exactly the equation we want to solve:

$$\underbrace{\frac{\hbar\omega}{2} \sigma_n}_{H} |F_j\rangle = \left( \frac{\hbar\omega}{2} F_j \right) |F_j\rangle$$

Multiplying the equation by a constant doesn't change the eigenvectors: they still are the only solutions, but the associated eigenvalues are now different:

$$\lambda_1 = \frac{\hbar\omega}{2}; \quad |\lambda_1\rangle = \begin{pmatrix} \cos(\theta/2) \\ \exp(i\phi) \sin(\theta/2) \end{pmatrix}$$

$$\lambda_2 = -\frac{\hbar\omega}{2}; \quad |\lambda_2\rangle = \begin{pmatrix} -\sin(\theta/2) \\ \exp(i\phi) \cos(\theta/2) \end{pmatrix}$$

<sup>1</sup>That's quite a fancy name for describing the eigenvectors of an operator, by comparison with the "iconic" Schrödinger equation...